

[HarvardScience](#) ▶ [Health & Medicine](#) ▶

Doctors can feel their patients' pain

Brain scans show physicians can feel pain, pain relief

By **Bonnie Prescott**

Beth Israel Deaconess Medical Center Communications

Tuesday, January 29, 2013




Image courtesy of Karin Jensen

"Now, for the first time, we've shown that caring for patients encompasses a unique neurobiology in physicians. Our ultimate goal is to transform the 'art of medicine' into the 'science of care,'" said the study's senior author, Ted Kaptchuk, an associate professor of medicine at Harvard Medical School. The image shows the regions in physicians' brains that were activated while the doctors were treating patients.

A patient's relationship with his or her doctor has long been considered an important component of healing. Now, in a novel investigation in which physicians underwent brain scans while they believed they were treating patients, researchers have provided the first scientific evidence indicating that doctors truly can feel their patients' pain — and can also experience their relief following treatment.

Led by researchers at Harvard-affiliated [Massachusetts General Hospital \(MGH\)](#) and the [Program in Placebo Studies and Therapeutic Encounter \(PiPS\)](#) at [Beth Israel Deaconess Medical Center \(BIDMC\)/Harvard Medical School \(HMS\)](#), the new findings, which appear online today in [Molecular Psychiatry](#), help to illuminate one of the more intangible aspects of health care — the doctor/patient relationship.

"Our findings showed that the same brain regions that have previously been shown to be activated when patients receive placebo therapies are similarly activated in the brains of doctors when they administer what they think are effective treatments," explained the first author, [Karin Jensen](#), an investigator in the Department of Psychiatry and Martinos Center for Biological Imaging at MGH and a member of the PiPS. Notably, she added, the findings also showed that the physicians who reported greater ability to take things from the patients' perspective, that is, to empathize with their feelings, experienced higher satisfaction during patients' treatments, as reflected in the brain scans.

"By demonstrating that caring for patients involves a complex set of brain events, including deep understanding of the patients' facial and body expressions, possibly in combination with the physician's own expectations of relief and feelings of reward, we have been able to elucidate the neurobiology underlying caregiving," added the senior author, [Ted Kaptchuk](#), director of the PiPS and an associate professor of medicine at HMS. "Our findings provide early evidence of the importance of interacting brain networks between patients and caregivers and acknowledge the doctor/patient relationship as a valued component of health care, alongside medications and procedures."

Previous investigations have demonstrated that a brain region associated with pain relief (the right ventrolateral prefrontal cortex, or VLPFC) and a region associated with reward (the rostral anterior cingulate cortex, or rACC) are activated when patients experience the placebo effect, which occurs when patients show improvement from treatments that contain no active ingredients.

Although behavioral research has suggested that physicians' expectations influence patients' clinical outcomes and help determine their placebo responses, until now little effort has been directed to understanding the biology underlying the physician component of the clinical relationship. Jensen and her colleagues hypothesized that the same brain regions that are activated during patients' placebo responses would similarly be activated in the brains of physicians as they treated patients. They also hypothesized that a physician's perspective-taking skills would influence the outcomes.

To test these hypotheses, the scientists developed a unique equipment arrangement that would enable them to conduct functional magnetic resonance imaging (fMRI) of the physicians' brains while the doctors had face-to-face interactions with patients, including observing patients as they underwent pain treatments.

The experiment included 18 physicians and two women who played the role of "patients" and followed a rehearsed script. The experiment called for the participating physicians to administer pain relief with what they thought was a pain-relieving electronic device, but which was actually a non-active "sham" device.

To ensure that the physicians believed that the sham device really worked, the investigators first administered a dose of "heat pain" to the physicians' forearms to gauge their pain threshold, and then "treated" them with the fake machine. During the treatments, the investigators reduced the heat stimulation, to demonstrate to the participants that the therapy worked. The physicians underwent fMRI scans while they experienced the painful heat stimulation so that investigators could see exactly which brain regions were activated during first-person perception of pain.

In the second portion of the experiment, each physician was introduced to a patient and asked to perform a standardized clinical examination. (The clinical exam was performed to establish a realistic rapport between the physician and patient before fMRI scanning took place, and was comparable to a standard U.S. doctor's appointment.) At that point the physician also answered a questionnaire, the Interpersonal Reactivity Index, used to measure the participant's self-reported perspective-taking skills.

During the third step, "The physician went inside the scanner and was equipped with a remote control that could activate the 'analgesic device' when prompted," Jensen explained. Mirrors inside the scanner enabled physicians to maintain eye contact with the patient, who was seated on a chair next to the scanner's bed and hooked up to both the thermal pain stimulator and the pain-relieving device.

Then, in a randomized order, physicians were instructed to either treat a patient's pain or to press a control button that provided no relief. When physicians were told not to activate pain relief, the "patient's" face registered pain while the physicians watched. When the physicians were instructed to treat the patients' pain, they saw that the subjects' faces were neutral and relaxed, the result of pain relief. During these doctor-patient interactions, fMRI scans measured the doctors' brain activations.

As predicted, the authors found that while treating patients, the physicians activated the right VLPFC region of the brain, a region previously implicated in the placebo response. Furthermore, Jensen said, the physicians' ability to take the patients' viewpoints correlated to brain activations and subjective ratings; physicians who reported high perspective-taking skills were more likely to show activation in the rACC brain region, which is associated with reward.

"We already know that the physician-patient relationship provides solace and can even relieve many symptoms," said Kaptechuk. "Now, for the first time, we've shown that caring for patients encompasses a unique neurobiology in physicians. Our ultimate goal is to transform the 'art of medicine' into the 'science of care,' and this research is an important first step in this process as we continue investigations to find out how patient-clinician interactions can lead to measurable clinical outcomes in patients."

[Tweet](#)

© 2013 The President and Fellows of Harvard College